

Sedum forsterianum

Oracle

Culture guide

Uses:

Pot, Rockery, dry and heat tolerant, roof garden, complementary

Exposure:

Sun

Garden height:

6" / 15 cm

Crop time:

8-10 months

Sow time:

January-March for green pots from April to August; June-August for flowering in pots the following year

Sowing method:

3-5 seeds per plug, sowing directly into final pot is recommended

Germination:

Germinates in 10-14 days at 65-72 °F (18-22 °C). Light improves germination and uniformity of emergence. Avoid excessive soil moisture.

Growing On:

Transplant plugs after 7 weeks. Once plants have rooted out, begin reducing soil moisture levels and lower temperatures to 55-62 °F (13-16 °C). Warmer temperatures will promote softer, more rapid growth. Feed at 100-150 ppm nitrogen in a well-balanced mix. Vernalization is required for flower initiation in every variety except 'Spirit'. However, most varieties are grown for their foliage characteristics rather than for their flowers

Media:

Use a well-drained, growing substrate with 0-15 % clay, 0-15 % parts (e.g. bark, wood fibres, perlite, sand), 1-1,5 kg/m³ complete balanced fertilizer, 1-2 kg/m³ slow release fertilizer (3-9 months), iron-chelate, micronutrien

Temperature:

Grow at 10-18 °C or outdoors. In winter indoors frost free at 3-5 °C or outdoors. Outdoor fleece cover needed. For wintering the root development should be very good. In spring the plants start to grow for 10-12 weeks at 15-18 °C. Cold temperatures at 10-12 °C will increase the cultivation time. A chilling period (vernalization) is required for flower initiation.

Fertilization:

Low-moderate fertilization levels are required.

Fertilize the crop weekly with 80-100 ppm nitrogen (at 2 kg/m³ slow release fertilizer in substrate), using complete balanced fertilizer.

Avoid high ammonium and high nitrogen levels.

Very high nitrogen levels in substrate cause shoot stretching and the shoots fall apart. Don't fertilize after mid September. In spring fertilize 80-100

ppm nitrogen of a complete balanced fertilizer.

Prevent magnesium deficiency by applying magnesium sulphate (0,05 %) 1-2 times and in case of iron deficiency (above pH 6.0) apply iron-chelate for 1-2 times.