

# Lavandula angustifolia Munstead Variety

Item no.: O9990/T

## Height

40cm

## Exposure

Sun

## Seed form

Raw Seed, ApeX

## Uses

Bedding, Cutflower, Landscape

## Culture guide

### Usage

Plants for border and open positions, ornamental leaf plant, aromatic plant, medicinal plant, pot and container plants, plants attractive for bees, extensive roof planting

### Sow time

January-March for green pots, June-August for flowering in pots the following year

### Sowing method

1-2 seeds per plug

### Germination

Germinates in 14-20 days at 65-72 °F (18-22 °C). Light is required for germination. Treated ApeX seed increases germination percentages over that of raw seed.

## **Growing on**

Transplant plugs after 9 weeks. Grow on at 55-65 °F (12-18 °C) in an alkaline medium. Vernalization is required for flower initiation. After vernalization, begin forcing plants at temperatures of 60-65 °F (15-18 °C) for 6-7 weeks under long days to promote flowering.

## **Media**

Use a well-drained, growing perennial substrate with 0-15 % clay, 0-15 % parts (e.g. perlite, sand, bark), 1-2 kg/m<sup>3</sup> complete balanced fertilizer, 0-2 kg/m<sup>3</sup> slow release fertilizer (3-9 months), iron-chelate, micronutrients, pH: 5.8-7.0.

## **Temperature**

Grow at 12-21 °C or outdoors. In winter indoors frost free at 3-5 °C or outdoors. Outdoor fleece cover needed. In spring the plants start to grow for 6-8 weeks at 18-21 °C. Cold temperatures at 12-15 °C will increase the cultivation time by 3-4 weeks. A chilling period (vernalization) for flower initiation for at least 15 weeks at 3-5 °C is required. A forcing in spring with warm temperatures need high light intensity and decreases the cultivation time.

---

Stage I Starts with the radicle breaking through the testa. The roots are touching the medium. Ends with fully developed cotyledons.

Stage II Starts from fully developed cotyledons. Ends with the fully developed true leaf or true leaf pair.

Stage III Starts from the fully developed true leaf or true leaf pair and ends with 80% of the young plants being marketable.

Stage IV All young plants are ready for sale and in the process of being hardened off. This stage lasts about 7 days.

The cultural recommendations are based on results from trials conducted under Central

European conditions. Different conditions in other parts of the world may lead to deviations in results achieved.