

Astilbe arendsii

Astary®

Mix (incl. Pink)

Item no.: AA0199P



- Fastrax perennial: First year flowering without vernalization
- Compact plant habit with single colors
- Perfect for pot production

| | |
|-----------------------|-----------------------|
| Crop Time | Spring: 18 - 20 weeks |
| Height ♂ | 10 " / 25 cm |
| Exposure | Sun - Shade |
| Seed Form | BeGreen Pelleting |
| Heat Zone | 8-2 |
| Hardiness Zone | 4-8 |
| Product Use | Bedding, Landscape |

TECHNICAL GUIDE

Astilbe arendsii Astary®

Usage

pot plant, border plant

Sow time

February – April

Sowing method

2-3 seeds per plug

Germination

Sow 2-3 seeds per cell in well-drained media with good moisture holding capacity. Maintain high but not saturated soil moisture levels during plug production. Germinates at 68-72 °F (20-22 °C) in 28 days. During Stage II lower temperatures to 65-70 °F (18-21 °C). Plugs should finish in 10-12 weeks. DO NOT VERNALIZE PLUGS.

Growing on

Grow on 65-70 °F (18-21 °C) - DO NOT VERNALIZE PLUGS

Media

Use a friable, coarse-fibered perennial media containing 10-20 % clay, 0-15 % organic matter (such as wood chips or bark humus), a multiple balanced fertilizer at 1-1.5 kg per m³, Radigen at 100 g/m³, and a coated time release fertilizer (4-5 months) at 3-4 kg; pH value: 5.5-6.0

Temperature

15-18 °C. Non-frost-free overwintering: Cover the plants with a non-woven cloth to protect against direct sunlight. Frost-free overwintering: 3-5 °C

Fertilization

Heavy feeder. Apply 150-200 mg N per l of a balanced multiple fertilizer containing a high percentage of nitrate nitrogen. Ensure a good potassium supply. Avoid overfertilization with nitrogen. Use 1-2 applications of sequestered iron to avoid iron deficiency.

Stage I Starts with the radicle breaking through the testa. The roots are touching the medium. Ends with fully developed cotyledons.

Stage II Starts from fully developed cotyledons. Ends with the fully developed true leaf or true leaf pair.

Stage III Starts from the fully developed true leaf or true leaf pair and ends with 80% of the young plants being marketable.

Stage IV All young plants are ready for sale and in the process of being hardened off. This stage lasts about 7 days.

The cultural recommendations are based on results from trials conducted under Central European conditions. Different conditions in other parts of the world may lead to deviations in results achieved.

COLORS OF THE SERIES

Astilbe arendsii Astary®



White
AA0103P



Rose
AA0102P



Mix (incl. Pink)
AA0199P