

Bellis perennis

Speedstar® Plus

Mix

Item no.: BP0299P



- Ideal for spring and fall production - perfect partner to Pansies
- Cultivation under cooler temperatures
- Early flowering, short crop time
- Compact plant habit, no PGRs required
- No vernalization required

| | |
|---------------------------------------------------------------------------------------------------|-----------------------------------------------|
| Crop Time | Spring: 17 - 20 weeks , Autumn: 10 - 14 weeks |
| Height  | 5 " / 13 cm |
| Exposure | Sun - Partial shade |
| Seed Form | Pelleted Seed |
| Best Uses | Bedding, Landscape, Pot Plant |

CULTURE GUIDE

Bellis perennis Speedstar® Plus

Usage

Flowering without vernalisation in fall and spring; Pot, mixed containers, borders or flower beds

Sow time

July-August for fall sales; September-November for spring sales

Sowing method

1 or 2 pellets per plug

Germination

7-14 days at 68 °F (20 °C), do not cover seed. After germination, gradually reduce temperatures to 60-68 °F (15-20 °C) while maintaining adequate ventilation.

Growing on

Grow on at temperatures of 65-68 °F (18-20 °C) for fall production and 45-50 °F (7-10 °C) for spring production. Plants will flower in approximately 8-10 weeks. Sow in July for fall flowering; or October for early spring flowering. Produce both fall and spring Bellis crops under conditions similar to pansy production.

Media

Use a well drained growing substrate, pH: 5.5-6.3

Temperature

16-18 °C for fall and early spring sales or frost free to 10 °C for later spring sales.

Fertilization

Moderate fertilization levels are required

Stage I Starts with the radicle breaking through the testa. The roots are touching the medium. Ends with fully developed cotyledons.

Stage II Starts from fully developed cotyledons. Ends with the fully developed true leaf or true leaf pair.

Stage III Starts from the fully developed true leaf or true leaf pair and ends with 80% of the young plants being marketable.

Stage IV All young plants are ready for sale and in the process of being hardened off. This stage lasts about 7 days.

The cultural recommendations are based on results from trials conducted under Central European conditions. Different conditions in other parts of the world may lead to deviations in results achieved.

COLORS OF THE SERIES

Bellis perennis Speedstar[®] Plus



Red
BP0203P



Rose
BP0202P



White
BP0204P



Mix
BP0299P