

# Callistephus chinensis Standy Light Pink

Item no.: F1141

## Crop Time

Spring: 14 - 15 weeks

## Height

75cm

## Exposure

Sun - Partial shade

## Seed Form

Raw Seed

## Best Uses

Bedding, Cutflower

## Culture guide

### Usage

Plants for the border, pot and container plants, plants for patio and window boxes, cut flowers

### Sow time

January for flowering in May (artificial long days); May for flowering in September (natural day length); July for flowering in January (artificial long days)

### Sowing method

3-5 seeds per plug, sowing in final pots is possible

### Germination

10-12 days at 68-70 °F (20-21 °C), cover seed lightly with vermiculite after sowing. To prevent premature flowering, keep plants vegetative with a four hour night interruption

## **Growing on**

Transplant plugs after 3-4 weeks. Grow on at 58-60 °F (14-16 °C). Cool temperatures promote a good plant habit, but may delay flowering.

## **Media**

Use a well-drained, growing substrate with 20-30 % clay, 1-1,5 kg/m<sup>3</sup> complete balanced fertilizer, 0-2 kg/m<sup>3</sup> slow release fertilizer (3-6 months), iron-chelate, micronutrients, pH: 6.0-6.5.

Use a sterile medium and sterile pots/containers as preventative measures against Fusarium and Verticillium. Field: sandy loamy soils with good drainage and moderate nutrition levels.

Standard fertilization: 80-100 g/m<sup>2</sup> slow release fertilizer. Disinfect the soils in greenhouse before planting.

## **Temperature**

Grow at 12-15 °C. Callistephus do not grow temperatures below 8 °C and do not tolerate frost.

The lower the temperature, the better the plant habit but onset of flowering will be a little delayed. Lower temperatures combined with short days cause the development of a leaf rosette and short stems.

---

Stage I Starts with the radicle breaking through the testa. The roots are touching the medium.

Ends with fully developed cotyledons.

Stage II Starts from fully developed cotyledons. Ends with the fully developed true leaf or true leaf pair.

Stage III Starts from the fully developed true leaf or true leaf pair and ends with 80% of the young plants being marketable.

Stage IV All young plants are ready for sale and in the process of being hardened off. This stage lasts about 7 days.

The cultural recommendations are based on results from trials conducted under Central European conditions. Different conditions in other parts of the world may lead to deviations in results achieved.