Carex comans

Zora

Item no.: CC5001R



- Finely textured ornamental grass
- Extraordinary, brownish color
- Versatile use in mixed containers or beds

Crop Time	Spring: 16 - 18 weeks
Height ⌀	14 " / 35 cm
Width ⌀	8 " / 20 cm
Exposure	Sun - Partial shade
Seed Form	Raw
Product Use	Bed, Landscape
Foliage Color	Brown

TECHNICAL GUIDE

Carex comans Zora

Plug Culture
Germination: Optimum conditions for seedling development, beginning on the day of sowing until radicle emergence. Expect germination in 7-14 days.

Cover: No cover is required.

Sowing method: 4-6 seeds per plug.

Media: pH 5.8-6.2; EC <1.0

Temperature: 20-22 °C (68-72 °F) until radicle emergence, then reduce to 18-20 °C (64-68 °F) during day and night.

Moisture: Begin with a wet (4) for the first days of germination. Alternate between moisture levels moist (3) and medium (2) to maintain a good wet to dry cycle. Avoid overwatering.

Light: Supplemental lighting is optional in the plug stage. If desired, provide light levels up to 35,000 lx.

Fertilizer: Maintain an EC <1.0. Use a nitrate-based fertilizer with low phosphorus. Feeding should start <100 ppm N and gradually work up to 100-175 ppm N.

Growth Regulators: Not required.

Growing On
Media: pH 5.5-6.2; EC 1.0-1.2

Temperature: 15-18 °C (58-64 °F) during day and night.

Moisture: Alternate between moisture levels moist (3) and medium (2) to maintain a good wet to dry cycle. Avoid overwatering.

Light: Lighting is not required.

Fertilizer: Feed moderate with 150-200 ppm N using a nitrate-based fertilizer with low phosphorus. Avoid overfeeding.

Growth Regulators: Not required.

Pests & Diseases: Aphids, root rot.

Plug & Finished Crop Time
Plug Crop Time

288 tray: 5-7 weeks

Finished Crop Time

10-12 cm (4-5") pots (1*): 9-10 weeks

15-19 cm (6-7") pots (3*): 11-12 weeks

Moisture Codes

Saturated (5) Water is easily observed when finger is pressed on cell. Water moves freely from the top of the plug to the bottom.

Wet (4)	Media looks black and is not glistening. The media feels wet to the touch but there is very little water movement.
Moist (3)	Water is not easily visible. When finger is pressed on the cell there is very little movement from top to bottom.
Medium (2)	Media is not black, but now looks medium brown. There is no water movement when pressed with finger.
Dry (1)	Media has changed color to a very light brown and is dry to the touch.

All information in our technical guide is based on our own trials and would therefore be as guideline only. Detailed cultivation aspects vary depending on climate, location, time of year and environmental conditions. Benary expressly disclaims any responsibility for the content of such data/information and makes no representation or warranty for the cultivation of any products listed. It is recommended that growers conduct a trial of products under their own conditions.

COLORS OF THE SERIES

Carex comans Zora



CC5001R