

# Celosia cristata Brainiac Robo Red

Item no.: H4181

## **Crop Time**

Spring: 10 - 11 weeks

## **Height**

15cm

## **Exposure**

Sun

## **Seed Form**

Raw Seed

## **Best Uses**

Pot Plant

## **Culture guide**

### **Usage**

Mass plantings, specimen plant and mixed containers

### **Sow time**

January-June

### **Sowing method**

1 seed per plug

### **Germination**

Stage 1: 7-10 days at 25 °C; Stage II-IV: 11-20 days at 18-21 °C day temperature and 15-18 °C night temperature. Use media with low soluble salt levels and pH: 5.5-6.0. Requires light for germination. Cover seed lightly with vermiculite after sowing. Keep soil slightly moist but not wet.

## Media

Use a well-drained, growing substrate with 15-30 % clay, 1-1,5 kg/m<sup>3</sup> complete balanced fertilizer, iron-chelate, micronutrients, pH: 5.5-6.0.

## Temperature

Grow at 18-20 °C. At the end of the cultivation time decrease the temperature to 16 °C. Avoid temperatures below 16 °C, because Celosia needs warm temperatures to grow.

## Tip

Cultivate plants firstly slightly moist and later dry. Avoid an inhibited and stressful cultivation (in particular low temperatures or nutrition deficiency), because the flower initiation may start before plants have reached sufficient size and thus the plant quality decreases.

---

Stage I Starts with the radicle breaking through the testa. The roots are touching the medium. Ends with fully developed cotyledons.

Stage II Starts from fully developed cotyledons. Ends with the fully developed true leaf or true leaf pair.

Stage III Starts from the fully developed true leaf or true leaf pair and ends with 80% of the young plants being marketable.

Stage IV All young plants are ready for sale and in the process of being hardened off. This stage lasts about 7 days.

The cultural recommendations are based on results from trials conducted under Central European conditions. Different conditions in other parts of the world may lead to deviations in results achieved.