



1. [Home](#)

Campanula carpatica

Pearl

- Significantly earlier – reduced crop time
- Great impulse item or hardy perennial
- Narrow flowering window

[Bookmark](#)

[Recommend](#)

[Print](#)

Crop Time

Spring: 17 - 20 weeks

Height

7 ? / 18 cm

Exposure

Sun - Partial shade

Seed Form

Raw Seed, Pelleted Seed

Heat Zone

9 - 5

Hardiness Zone

12 - 8

Best Uses

Bedding, Landscape, Rockery

Culture guide

Usage

Pots, Hanging Baskets, Mixed Containers and Landscape

Sowing method

Raw seed: sow 6-8 seeds / plug; Pelleted seed: sow 2-3 pellets / plug

Germination

Optimum conditions for seedling development, beginning on the day of sowing until radical emergence. Expect radicle emergence in 10-12 days

Temperature

68-72 ?F (20-22 ?C) day temperature, 61 ?F (18 ?C) night temperature. The fluctuation between day and night temperatures will aid in the germination. By day 14 the temperature can

be maintained at 64-68 °F (17-20 °C).

Fertilization

Sensitive to high EC in the early stages so maintain and EC less than 0.5 for the first 10-14 days until germination is complete.

Stage I Starts with the radicle breaking through the testa. The roots are touching the medium. Ends with fully developed cotyledons.

Stage II Starts from fully developed cotyledons. Ends with the fully developed true leaf or true leaf pair.

Stage III Starts from the fully developed true leaf or true leaf pair and ends with 80% of the young plants being marketable.

Stage IV All young plants are ready for sale and in the process of being hardened off. This stage lasts about 7 days.

The cultural recommendations are based on results from trials conducted under Central European conditions. Different conditions in other parts of the world may lead to deviations in results achieved.

[Download](#)