

Begonia x hybrida F? Funky® Light Pink

Item no.: C8530P

Crop Time

Spring: 15 - 18 weeks

Height

25cm

Exposure

Sun - Shade

Seed Form

Pelleted Seed

Best Uses

Bedding, Hanging basket

Culture guide

Usage

Pots, Hanging Baskets, Mixed Containers and Landscape

Sowing method

Sow 1-2 seeds or pellets per plug.

Germination

Optimum conditions for seedling development, beginning on the day of sowing until radicle emergence. Expect radicle emergence in 7–10 days. Germination should be complete at 10 days unless optimum conditions are not provided.



Temperature

Plug Culture:

72–74 °F (22-23 ?C) until radicle emergence. Higher temperatures, exceeding 80 °F (27 °C) will inhibit germination. Upon radicle emergence, on day 10-14 reduce the temperature to 68–70 °F (20–21 °C) until cotyledon expansion has occured. On day 21 the temperature can be reduced further to 68 °F (20 °C). For irrigation use tempered water (above 64 °F / 18 °C) only. Growing on:

After transplanting 65–68 °F (18–20 °C) nights for the first 14 days or until the roots reach the bottom of the container. Thereafter temperatures may be lowered to 62–64 °F (17–18 °C). An ADT (average daily temperature of 67 ?F (19 ? C) will give the fastest finished crop. Temperatures below 57 ?F (14 ?C) will result in tuber formation and a delay of the crop. A DIFF of 2-3 ?F will result in a more compact crop requiring little to no growth regulators.

Tip

Begonia Funky® is a showstopper with single, double and semi double flowers on the same plant. Reliable high germination rates and easy to grow & ship make this one a no-brainer!

Stage I Starts with the radicle breaking through the testa. The roots are touching the medium. Ends with fully developed cotyledons.

Stage II Starts from fully developed cotyledons. Ends with the fully developed true leaf or true leaf pair.

Stage III Starts from the fully developed true leaf or true leaf pair and ends with 80% of the young plants being marketable.

Stage IV All young plants are ready for sale and in the process of being hardened off. This stage lasts about 7 days.

The cultural recommendations are based on results from trials conducted under Central European conditions. Different conditions in other parts of the world may lead to deviations in results achieved.