





1. Home

Delphinium grandiflorum

Summer

- FastraX perennial: First year flowering without vernalization
- Compact plant habit
- Perfect as mixture in containers

Bookmark

Recommend



Print

Crop	Time
------	------

Spring: 16 - 20 weeks

Height?

14?/35 cm

Exposure

Sun - Partial shade

Seed Form

Raw Seed

Heat Zone

6-1

Hardiness Zone

3b-7b

Best Uses

Bedding, Cutflower, Landscape

Culture guide

Usage

Perennial, first year flowering. Bedding and border plants, pot and mixed containers.

Sow time

December-May

Sowing method

2-3 seeds per plug

Germination



14-18 days at 68-72 °F (20-22 °C). Cover seed lightly and maintain constant humidity levels during Stage I.

Growing on

Transplant plugs 6-8 weeks after sowing into 3.5" (9 cm) to 1 quart (1 liter) containers. For larger containers, use 3 plants per pot. Avoid planting plugs too deep. Provide good ventilation. Begin fertilizing at 200 ppm nitrate nitrogen in a 2-1-2 ratio fertilizer mix. Avoid soluble salt buildups. Finish at temperatures of 45-75 °F (7-24 °C). Temperatures above 85 °F (29 °C) can cause flower abortion. Temperatures below 55 °F (13 °C) under short days will cause plants to form rosettes and will delay flowering. At lower temperatures, use extended day length or night interruption with incandescent lights to avoid rosette formation.

Media

Use a well-drained media, pH: 5.8-6.5.

Temperature

Grow at 13-16 °C or outdoors.

Fertilization

Moderate fertilization levels are required.

Stage I Starts with the radicle breaking through the testa. The roots are touching the medium. Ends with fully developed cotyledons.

Stage II Starts from fully developed cotyledons. Ends with the fully developed true leaf or true leaf pair.

Stage III Starts from the fully developed true leaf or true leaf pair and ends with 80% of the young plants being marketable.

Stage IV All young plants are ready for sale and in the process of being hardened off. This stage



lasts about 7 days.

The cultural recommendations are based on results from trials conducted under Central European conditions. Different conditions in other parts of the world may lead to deviations in results achieved.

Download