

Portulaca grandiflora F? Sundial Caliente

Mix

Item no.: U1701/U

Crop Time

Spring: 8 - 10 weeks

Height

13cm

Exposure

Sun

Seed Form

Raw Seed, Multipelleted Seed

Best Uses

Bedding, Landscape, Rockery

Culture guide

Usage

Mixed containers, bedding, landscaping

Sowing method

2 seeds per plug, 1 multi-seed pellet per plug

Germination

Stage I: 2-3 days at 70-78 °F (22-26 °C). Sow in media with very low soluble salt levels and pH 5.5-6.2. Light is not essential, but beneficial for germination. Keep soil moist but not wet. Keep

relative humidity between 90-95%. At the end of stage I the radicle will have emerged and the cotyledons begin to unfurl.

Growing on

4-5 weeks after sowing (288-tray) transplant 1 plant into 10-11 cm (4-5") pots or packs

Media

Use a well-drained substrate with pH 5.5 -6.2 and 1-1.5kg/m³ fertilizer

Temperature

18 °C-22 °C (65-72 °F). Temperatures below 18 °C (65 °F) will inhibit induction and result in poor or no vegetative growth.

Stage I Starts with the radicle breaking through the testa. The roots are touching the medium. Ends with fully developed cotyledons.

Stage II Starts from fully developed cotyledons. Ends with the fully developed true leaf or true leaf pair.

Stage III Starts from the fully developed true leaf or true leaf pair and ends with 80% of the young plants being marketable.

Stage IV All young plants are ready for sale and in the process of being hardened off. This stage lasts about 7 days.

The cultural recommendations are based on results from trials conducted under Central European conditions. Different conditions in other parts of the world may lead to deviations in results achieved.