

Lobelia fulgens Queen Victoria

Item no.: R3240P

Crop Time

Spring: 18 - 20 weeks

Height

75cm

Exposure

Sun - Partial shade

Seed Form

ApeX Pelleted

Hardiness Zone

6-10

Best Uses

Bedding, Cutflower, Landscape

Culture guide

Usage

Perennial for border and cottage garden, container plant, bedding, cut flower plants

Sow time

January-March for flowering in pots from July onwards, June-August for flowering in pots the following year, April-May for cut flower production outdoors

Sowing method

1 seed per plug for short day conditions

3-5 seeds per plug for long day condition, because at long day the plants do not form a leaf rosette.

Germination

Germinates in 14-20 days at 65-72 °F (18-22 °C). Light is required for germination.

Growing on

Transplant plugs after 8-9 weeks. Grow on at 60-65 °F (15-18 °C). Vernalization is not required for flower initiation. After transplant, maintain temperatures of 60-65 °F (15-18 °C) for 7-8 weeks under long days.

Media

Use a well-drained, growing substrate with 15-30 % clay, 1-1,5 kg/m³ complete balanced fertilizer, 0-3 kg/m³ slow release fertilizer (3-9 months), iron-chelate, micronutrients, pH: 6.0-7.0. Field: humus, sandy humus soils with a good drainage. Standard fertilization: 60-80 g/m² of a slow release fertilizer.

Temperature

Grow at 15-18 °C, later decrease the temperature to 10-15 °C, when the root development in pots is very good. Cultivation outdoors is possible, too. In winter indoors frost free at 3-5 °C or outdoors. Outdoors fleece cover needed. *L. fulgens* are very sensitive to very strong frost temperatures. In spring the plants start to grow at 15-18 °C. Cold temperatures at 10-12 °C will increase the cultivation time. A chilling period (vernalization) for flower initiation is not required.

Stage I Starts with the radicle breaking through the testa. The roots are touching the medium. Ends with fully developed cotyledons.

Stage II Starts from fully developed cotyledons. Ends with the fully developed true leaf or true leaf pair.

Stage III Starts from the fully developed true leaf or true leaf pair and ends with 80% of the young plants being marketable.

Stage IV All young plants are ready for sale and in the process of being hardened off. This stage lasts about 7 days.

The cultural recommendations are based on results from trials conducted under Central European conditions. Different conditions in other parts of the world may lead to deviations in results achieved.