

Viola wittrockiana F? Inspire® Plus Red

Blotch

Item no.: X0315/E

Crop Time

Spring: 25 - 26 weeks , Autumn: 12 - 10 weeks

Height

18cm

Exposure

Sun

Seed Form

Raw Seed, BeGreen Priming

Best Uses

Bedding, Landscape

Culture guide

Usage

Packs, Pots, Mixed Containers and Landscape/Mass plantings

Sowing method

1 seed per plug

Germination

Optimum conditions for seedling development, beginning on the day of sowing until radicle emergence. Expect radicle emergence in 2-4 days.

Temperature

65-68 °F (18-20 °C) until radicle emergence, then lower the temperature gradually to 62-65 °F (17-18 °C).

Stage I Starts with the radicle breaking through the testa. The roots are touching the medium. Ends with fully developed cotyledons.

Stage II Starts from fully developed cotyledons. Ends with the fully developed true leaf or true leaf pair.

Stage III Starts from the fully developed true leaf or true leaf pair and ends with 80% of the young plants being marketable.

Stage IV All young plants are ready for sale and in the process of being hardened off. This stage lasts about 7 days.

The cultural recommendations are based on results from trials conducted under Central European conditions. Different conditions in other parts of the world may lead to deviations in results achieved.