



1. [Home](#)

Viola wittrockiana F?

Inspire® DeluXXe

Indulge your Customers in Luxury!

- Leading series in extra large-flowered Pansies
- Huge, early and long lasting flowers
- Unmatched uniformity in habit and timing
- Superior outdoor performance
- Perfect for late summer / early fall production



[Bookmark](#)

[Recommend](#)

[Print](#)

Crop Time

Autumn: 10 - 11 weeks

Height ?

7 ? / 18 cm

Width ?

7 ? / 18 cm

Exposure

Sun

Seed Form

Raw Seed, BeGreen Priming

Best Uses

Bedding, Landscape

Culture guide

Usage

Packs, Pots, Mixed Containers and Landscape/Mass plantings

Sowing method

1 seed per plug

Germination

Optimum conditions for seedling development, beginning on the day of sowing until radicle emergence. Expect radicle emergence in 2-4 days.

Temperature

65-68 °F (18-20 °F) until radicle emergence, then lower the temperature gradually to 62-65 °F (17-18 °C).

Fertilization

Begin feeding early using a calcium based fertilizer at lower rates to keep the an adequate amount of calcium and nitrogen supplied to the seedlings. On days 5-7 begin feeding with a calcium based fertilizer (14-2-14, 13-2-13, 15-5-15, 17-5-17) at 50-60 ppm. Maintain the EC between 0.5 and 0.75. Keep phosphorous levels between 6-8 ppm and boron supplied at 0.5 ppm.

Stage I Starts with the radicle breaking through the testa. The roots are touching the medium. Ends with fully developed cotyledons.

Stage II Starts from fully developed cotyledons. Ends with the fully developed true leaf or true leaf pair.

Stage III Starts from the fully developed true leaf or true leaf pair and ends with 80% of the young plants being marketable.

Stage IV All young plants are ready for sale and in the process of being hardened off. This stage lasts about 7 days.

The cultural recommendations are based on results from trials conducted under Central European conditions. Different conditions in other parts of the world may lead to deviations in results achieved.

[Download](#)