

Myosotis sylvatica

Bellamy

Blue

Item no.: MS0401R



Benary's Popular & Well-Branched Myosotis Series

- Easy to grow, no cold required
- For early spring and fall production
- Well-branched, uniform plant habit
- Very uniform flowering across the series

Crop Time	Spring: 20 - 24 weeks , Autumn: 10 - 12 weeks
Height ♂	9 " / 23 cm
Width ♂	7 " / 18 cm
Exposure	Sun - Partial shade
Seed Form	Raw Seed
Hardiness Zone	5a-8a
Product Use	Pots, Landscape, Mixed Containers
Family, Origin	Boraginaceae, Europe
Minimum Germ. Rate	85%

TECHNICAL GUIDE

Myosotis sylvatica Bellamy

Usage

Pots, Mixed Containers and Landscape

Sowing method

1 seed per plug.

Germination

Maintain optimal conditions for seedling development, should begin on the day of sowing until root emergence. Expect root emergence in 3-6 days from sowing.

Temperature

Plug culture: 20-22 °C (68-72 °F) until radicle emergence. Afterwards, ensure 18-20 °C (64-68 °F) during night and day.

When the roots reach the bottom of the cell, the temperature can be lowered to 16-18 °C (60-64 °F).

Growing on: 16-20 °C (60-68 °F) days, 10-13 °C (50-56 °F) nights after transplanting. Afterwards, a frost-free production with a minimum temperature of 3-5 °C (38-41 °F) at night is recommended. Cooler temperatures promote a stable, compact and well-branched plant habit.

Fertilization

Plug culture: Maintain low pH level to avoid chlorosis. Begin feeding once germination is complete with 100-175 ppm N and maintain an EC of less than 1.2.

Growing on: Start feeding one week after transplanting. Apply regularly 175 to 225 ppm N by using predominantly nitrate-based fertilizer with low phosphorus. If needed, alternate with a balanced ammonium and nitrate-based fertilizer to encourage growth and balance the media pH. Maintain low pH level below 5.8 to avoid chlorosis.



COLORS OF THE SERIES

Myosotis sylvatica Bellamy



White
MS0403R



Pink
MS0402R



Blue
MS0401R



Light Blue
MS0404R



Mix
MS0499R