


Callistephus chinensis


Benary's Princess

Bright Rose

Item no.: CC0306R



- Tolerant of challenging growing conditions
- Large, dense filled flowers
- Strong stems as perfect cut flower
- Clear colors with bright yellow centers
- White is slightly taller than other colors

Crop Time	Spring: 14 - 15 weeks
Height 	28 " / 70 cm
Exposure	Sun - Partial shade
Seed Form	Raw Seed
Best Uses	Bedding, Cutflower

CULTURE GUIDE

Callistephus chinensis Benary's Princess

Usage

Plants for the border, pot and container plants, plants for patio and window boxes, cut flowers

Sow time

January for flowering in May (artificial long days); May for flowering in September (natural day length); July for flowering in January (artificial long days)

Sowing method

3-5 seeds per plug, sowing in final pots is possible

Germination

10-12 days at 68-70 °F (20-21 °C), cover seed lightly with vermiculite after sowing. To prevent premature flowering, keep plants vegetative with a four hour night interruption

Growing on

Transplant plugs after 3-4 weeks. Grow on at 58-60 °F (14-16 °C). Cool temperatures promote a good plant habit, but may delay flowering.

Media

Use a well-drained, growing substrate with 20-30 % clay, 1-1,5 kg/m³ complete balanced fertilizer, 0-2 kg/m³ slow release fertilizer (3-6 months), iron-chelate, micronutrients, pH: 6.0-6.5. Use a sterile medium and sterile pots/containers as preventative measures against *Fusarium* and *Verticillium*. Field: sandy loamy soils with good drainage and moderate nutrition levels. Standard fertilization: 80-100 g/m² slow release fertilizer. Disinfect the soils in greenhouse before planting.

Temperature

Grow at 12-15 °C. *Callistephus* do not grow temperatures below 8 °C and do not tolerate frost. The lower the temperature, the better the plant habit but onset of flowering will be a little delayed. Lower temperatures combined with short days cause the development of a leaf rosette and short stems.

Fertilization

Moderate-high fertilization levels are required. Fertilize the crop weekly with 100-150 ppm nitrogen (at 2 kg/m³ slow release fertilizer in substrate), using a potassium balanced fertilizer (N: K₂O-ratio: 1:1,5). Avoid high ammonium and high nitrogen levels. High nitrogen levels will cause smooth stems. Prevent magnesium deficiency by applying magnesium sulphate (0,025 %) 1-2 times and in case of iron deficiency apply iron-chelate for 1-2 times.

Field: N min soil value: approximately 150 g/m².

Stage I Starts with the radicle breaking through the testa. The roots are touching the medium. Ends with fully developed cotyledons.

Stage II Starts from fully developed cotyledons. Ends with the fully developed true leaf or true leaf pair.

Stage III Starts from the fully developed true leaf or true leaf pair and ends with 80% of the young plants being marketable.

Stage IV All young plants are ready for sale and in the process of being hardened off. This stage lasts about 7 days.

The cultural recommendations are based on results from trials conducted under Central European conditions. Different conditions in other parts of the world may lead to deviations in results achieved.

COLORS OF THE SERIES

Callistephus chinensis Benary's Princess



Blue
CC0301R



Bright Red
CC0307R



Bright Rose
CC0306R



Crimson
CC0304R



Deep Red
CC0309R



Light Blue
CC0303R



Salmon Rose
CC0305R



White
CC0308R



Yellow
CC0302R



Mix
CC0399R