

Kniphofia uvaria Flamenco

Item no.: N9381E

Crop Time

Spring: 26 - 28 weeks

Height

80cm

Exposure

Sun

Seed Form

BeGreen Priming

Hardiness Zone

5b-9b

Best Uses

Bedding, Cutflower, Landscape

Culture guide

Usage

Evergreen perennial for border and bedding, cut flowers

Sow time

January-May for flowering in pots from July onwards, June-August for flowering in pots the following year

Sowing method

2-3 seeds per plug, can be sown directly in final pots (4-6 seeds)

Germination

Germinates in 14-20 days at 65-72 °F (18-22 °C). Light is required for germination. During Stage II, maintain temperatures of 65-68 °F (18-20 °C). During Stages II and IV, gradually reduce temperature to 58-62 °F (14-16 °C).

Growing on

Transplant plugs after 8 weeks. Grow on at of 60 °F (15 °C) in a well-drained, humus, loamy medium. Vernalization is not required for flowering

Media

Use a well-drained, growing perennial substrate with 0-15 % clay, 0-15 % parts (e.g. bark, wood fibres), 1,5-2 kg/m³ complete balanced fertilizer, 2-3 kg/m³ slow release fertilizer (3-9 months), iron-chelate, micronutrients, pH: 5.5-6.2. Field: humus sandy soils, with a good drainage and the soil should warm up quickly. Standard fertilization: 60-80 g/m² of a slow release fertilizer.

Temperature

Grow at 15 °C or outdoors. In winter indoors frost free at 3-5 °C or outdoors. Outdoors a fleece cover or mulch cover with straw (20 cm) is needed. In spring the plants start to grow at 12-15 °C and long day. A chilling period for flower initiation is not required, but with a chilling period the flowering is more uniform and full. Kniphofia can also be made to flower 2-4 weeks earlier in the poly-tunnel.

Stage I Starts with the radicle breaking through the testa. The roots are touching the medium. Ends with fully developed cotyledons.

Stage II Starts from fully developed cotyledons. Ends with the fully developed true leaf or true leaf pair.

Stage III Starts from the fully developed true leaf or true leaf pair and ends with 80% of the

young plants being marketable.

Stage IV All young plants are ready for sale and in the process of being hardened off. This stage lasts about 7 days.

The cultural recommendations are based on results from trials conducted under Central European conditions. Different conditions in other parts of the world may lead to deviations in results achieved.