

# Campanula carpatica F? Pearl White

Item no.: H1040P

## Crop Time

Spring: 17 - 20 weeks

## Height

18cm

## Exposure

Sun - Partial shade

## Seed Form

Pelleted Seed

## Hardiness Zone

3a-8a

## Best Uses

Bedding, Landscape, Rockery

## Culture guide

### Usage

Pots, Hanging Baskets, Mixed Containers and Landscape

### Sowing method

Raw seed: sow 6-8 seeds / plug; Pelleted seed: sow 2-3 pellets / plug

### Germination

Optimum conditions for seedling development, beginning on the day of sowing until radical emergence. Expect radicle emergence in 10-12 days.

No cover is necessary.

## **Temperature**

68-72 °F (20-22 °C) day temperature, 61 °F (18 °C) night temperature. The fluctuation between day and night temperatures will aid in the germination. By day 14 the temperature can be maintained at 64-68 °F (17-20 °C).

---

Stage I Starts with the radicle breaking through the testa. The roots are touching the medium. Ends with fully developed cotyledons.

Stage II Starts from fully developed cotyledons. Ends with the fully developed true leaf or true leaf pair.

Stage III Starts from the fully developed true leaf or true leaf pair and ends with 80% of the young plants being marketable.

Stage IV All young plants are ready for sale and in the process of being hardened off. This stage lasts about 7 days.

The cultural recommendations are based on results from trials conducted under Central European conditions. Different conditions in other parts of the world may lead to deviations in results achieved.