

Limonium perezii



The Heat and Drought Tolerant Evergreen



- Violet-blue flowers on healthy, dark-green leaves
- Suitable for pots up to premium containers
- Optimal use in dry bouquets
- Attracts many beneficial insects

Technical Guide: Click here

All information in our technical guide is based on our own trials and would therefore be as guideline only. Detailed cultivation aspects vary depending on climate, location, time of year and environmental conditions. Benary expressly disclaims any responsibility for the content of such data/information and makes no representation or warranty for the cultivation of any products listed. It is recommended that growers conduct a trial of products under their own conditions.

Crop Time	Spring: 12 - 16 weeks
Height ∅	22 " / 55 cm
Width Ø	14 " / 35 cm
Exposure	Sun - Partial shade
Seed Form	Raw Seed
Product Use	Pots, Landscape, Cutflower, Containers



TECHNICAL GUIDE

Limonium perezii Indie

Usage

Pots, Mixed Containers and Landscape

Sowing method

1-3 seeds per plug, depending on the pot size.

Germination

Optimum conditions for seedling development, beginning on the day of sowing until radicle emergence. Expect radicle emergence in 7-10 days. A germination chamber promotes uniform germination

Temperature

Plug culture: 22-24 °C (72-76 °F) until radicle emergence. Afterwards, ensure 20 °C (68 °F) during night and day. When the roots reach the bottom of the cell, the temperature can be lowered to 16-18 °C (60-64 °F).

Growing on: 12-15 °C (54-58 °F) days and nights. Temperatures above 25 °C (77 °F) will inhibit the flower development and will increase the risk of infects with diseases. In addition, the leaves get bigger. Cooler temperatures promote stable flower stems and a compact, well-branched plant habit in pots. A frost-free outdoor production at 3-5 °C (38-40 °F) is also possible, but Limonium does not tolerate frost.

Stage I Starts with the radicle breaking through the testa. The roots are touching the medium. Ends with fully developed cotyledons. Stage II Starts from fully developed cotyledons. Ends with the fully developed true leaf or true leaf pair.

Stage III Starts from the fully developed true leaf or true leaf pair and ends with 80% of the young plants being marketable.

Stage IV All young plants are ready for sale and in the process of being hardened off. This stage lasts about 7 days.

The cultural recommendations are based on results from trials conducted under Central European conditions. Different conditions in other parts of the world may lead to deviations in results achieved.



COLORS OF THE SERIES

Limonium perezii Indie

