

Corynephorus canescens

Spiky

Item no.: CC5501R



Silvery-Blue Foliage

- Compact growth of blue-silvery foliage
- Drought tolerant, hardy grass
- Adds texture and contrast to landscape planting

Crop Time	Spring: 14 - 16 weeks
Height ∅	18 " / 45 cm
Width ∅	18 " / 45 cm
Exposure	Sun - Partial shade
Seed Form	Raw Seed
Hardiness Zone	5-9
Product Use	Landscape
Foliage Color	Green-Blue
Plume Color	White

TECHNICAL GUIDE

Corynephorus canescens Spiky

Sowing method

4-6 seeds per plug.

Germination

Optimum conditions for seedling development, beginning on the day of sowing until radicle emergence. Expect germination in 3-6 days.

Temperature

Plug culture: 20-22 °C (68-72 °F) until radicle emergence, then reduce to 18-20 °C (64-68 °F) during day and night.

Growing on: 15-20 °C (59-68 °F) during day and night. Cooler temperatures up to a minimum of 10 °C (50 °F) promote a compact, round plant habit in pots, but the crop time will increase significantly

Stage I Starts with the radicle breaking through the testa. The roots are touching the medium. Ends with fully developed cotyledons.

Stage II Starts from fully developed cotyledons. Ends with the fully developed true leaf or true leaf pair.

Stage III Starts from the fully developed true leaf or true leaf pair and ends with 80% of the young plants being marketable.

Stage IV All young plants are ready for sale and in the process of being hardened off. This stage lasts about 7 days.

The cultural recommendations are based on results from trials conducted under Central European conditions. Different conditions in other parts of the world may lead to deviations in results achieved.

COLORS OF THE SERIES

Corynephorus canescens Spiky



CC5501R