


Petunia x hybrida F₁

SUCCESS!® TR

Light Pink Morn

Item no.: PH0118P



- Trailing grandiflora Petunia
- Very uniform series in flower timing and habit
- Earliest trailing series on the market
- Unbeatable, intense colors and star types

Technical Guide: [Click here](#)

All information in our technical guide is based on our own trials and would therefore be as guideline only. Detailed cultivation aspects vary depending on climate, location, time of year and environmental conditions. Benary expressly disclaims any responsibility for the content of such data/information and makes no representation or warranty for the cultivation of any products listed. It is recommended that growers conduct a trial of products under their own conditions.

Crop Time	Spring: 11 - 13 weeks
Height ∅	11 " / 28 cm
Width ∅	26 " / 65 cm
Exposure	Sun - Partial shade
Seed Form	Pelleted Seed
Best Uses	Bedding, Hanging basket, Pot Plant, Landscape, Containers

CULTURE GUIDE

Petunia x hybrida F₁ SUCCESS!® TR

Usage

Pots, Hanging Baskets, Mixed Containers and Landscape

Sowing method

1 seed/pellet per plug.

Media

Plug culture: pH 5.5-5.8; keeping the pH below 6.0 will help to keep boron and iron available; EC 1.0 < or 0.75-1.0

Growing on: pH 5.5-5.8; EC 1.5-2.0

Temperature

Plug Culture: Maintain 22-24 °C (72-76 °F) until root emergence and then reduce to 20-21 °C (68-70 °F). The temperature can be lowered on approximately day 5. Once cotyledons have expanded lower temperature further to 18-20 °C (64-68 °F). Maintain 18-20 °C (64-68 °F) until day 26-28, then reduce the temperature to 15.5-18 °C (60-64 °F). Keep temperatures > 16 °C (60 °F) until ready to transplant. For the fastest finish maintain an ADT of 19.5 °C (67 °F). With these temperatures some additional growth regulators will need to be applied.

Growing on: After transplant maintain temperatures > 13 °C (56 °F) nights for the first 6 weeks to initiate flower bud development. The night temperatures can be lowered further to 10 °C (50 °F) to encourage basal branching and compactness. However, lower temperatures may also substantially decrease the number of flowers initiated. Growing at cooler temperatures will produce a higher quality plant. An ADT (average daily temperature) of 19 °C (66 °F) will give the fastest finished crop.

Fertilization

Plug culture: Maintain an EC < 1.0

Growing on: Fertilization: Feed at 100-200 ppm nitrogen. Under high light conditions, apply an ammonium-based feed (17-5-17). To prevent stretching under low light conditions apply a calcium-based feed (14-4-14). Under high light and long days an ammonium-based feed (20-10-20) is preferred.

Stage I Starts with the radicle breaking through the testa. The roots are touching the medium. Ends with fully developed cotyledons.

Stage II Starts from fully developed cotyledons. Ends with the fully developed true leaf or true leaf pair.

Stage III Starts from the fully developed true leaf or true leaf pair and ends with 80% of the young plants being marketable.

Stage IV All young plants are ready for sale and in the process of being hardened off. This stage lasts about 7 days.

The cultural recommendations are based on results from trials conducted under Central European conditions. Different conditions in other parts of the world may lead to deviations in results achieved.

COLORS OF THE SERIES

Petunia x hybrida F₁ SUCCESS!® TR



Blue
PH0102P



Burgundy
PH0103P



Light Pink Morn
PH0118P



Magenta Star
PH0115P



Pink
PH0104P



Pink Morn
PH0114P



Red
PH0105P



Rose
PH0106P



Rose Star
PH0116P



Salmon
PH0107P



Silver Vein
PH0101P



Violet
PH0109P



White
PH0110P



Maxi Mix
PH0199P